

Depolarization of Pulsar Radio Emission

Maxim Lyutikov

Canadian Institute for Theoretical Astrophysics, 60 St. George, Toronto, Ont, M5S 3H8, Canada

ABSTRACT

We show that intensity dependent depolarization of single pulses (e.g. Xiluoris et al. 1994) may be due to the nonlinear decay of the "upper" ordinary mode into an unpolarized extraordinary mode and a backward propagating wave. The decay occurs in the innermost parts of the pulsar magnetosphere for obliquely propagating O waves.

Subject headings: stars:pulsars-plasmas-waves-radiative transfer

1. Introduction

There is a marked anticorrelation between the intensity of a pulse and a degree of linear polarization (Manchester et al. 1975, Xilouris et al. 1994). Depolarization may be due to either a generic feature of the emission mechanism or wave propagation in the pulsar magnetosphere. In the latter case two mechanisms are possible - linear and nonlinear coupling. Linear coupling is due to the breakdown of the geometric optics approximation for polarization changes (Zheleznyakov 1996). The efficiency of linear coupling is independent of the wave amplitude. Nonlinear coupling is due to the wave-wave interaction processes. Its efficiency is proportional to the amplitudes or intensities of the interacting waves. The observed inverse dependence of the degree of linear polarization on intensity suggests that it is the nonlinear mechanism that is responsible for depolarization.

The pair plasma of pulsar magnetospheres supports three types of linearly polarized waves: transverse extraordinary (X) mode, and mixed Alfvén (A) and ordinary (O) modes. For a given wave vector the frequencies of the waves satisfy $\omega_O > \omega_X > \omega_A$, so that the possible induced decays are $O \rightarrow 2X$, $O \rightarrow X + A$, $X \rightarrow A + X$. In all cases the product waves propagate in approximately opposite directions.¹

Suppose that the emission mechanism generates waves of the ordinary (O) mode (some emission mechanisms do favor the O mode for the generation of radio emission). Initially the O mode may be strongly polarized. The preferred polarization of the emitted O mode may be related to the osculating plane of the curved magnetic field lines: the emission mechanism may generate waves polarized mostly in the osculating plane (e.g. curvature emission) or perpendicular to the this plane (Cherenkov-drift emission (Lyutikov et al. 1999)). As the emitted O mode propagates in the magnetosphere, it will experience induced decay into X modes and backward propagating waves. In the leading order of expansion in $1/\omega_B$ and for small angles of propagation with respect to the magnetic field, the decay of the O mode will produce two orthogonally polarized modes with *random polarization* with respect to the initial O mode (appendix A). The rate of this decay is proportional to the intensity of the wave (the rate of decay is proportional to the wave amplitude for the spectral narrow line, and to the wave intensity for the broad spectrum). The

¹Decay of X into two forward propagating X modes is suppressed by magnetic field - the matrix element $\sim 1/\omega_B^2$, compare with (Eq. A2).

product X modes can in turn decay into Alfvén waves. But Alfvén modes cannot escape from plasma and are damped by the Cherenkov interaction with plasma particles (Arons & Barnard 1986). Thus, the induced decay of $X \rightarrow A + X$ (with the X mode propagating backwards) is of no interest to us since it generates unobservable waves ².

The processes that will contribute to depolarization include decays (Fig. 1) $O \rightarrow 2X$ and $O \rightarrow X + A$ (in the latter case it is the Alfvén wave that is propagating backwards), and inverse processes of coalescence of backward propagating X or Alfvén mode with a forward propagating X mode. The inverse processes will be proportional to the product of the intensities of coalescing waves. In some case this may be an important process, but the convection of backward propagating waves from the interaction region will most probably reduce their intensity considerably making the coalescence ineffective. If inverse processes are ineffective, virtually all of the O mode may be converted into the X mode; the X modes may then be as intense as the initial O mode (backward propagating waves carry off only a small fraction of energy).

2. Induced decay of O mode

For the three wave processes to take place, the participating waves should satisfy resonance conditions (conservation of energy and momentum along the field)

$$\begin{aligned}\omega_o &= \omega_X + \omega_2 \\ \mathbf{k}_O &= \mathbf{k}_X + \mathbf{k}_2\end{aligned}\tag{1}$$

and more subtle conditions on the polarization determined by the matrix elements $A(\mathbf{k}_0, \mathbf{k}_X, \mathbf{k}_2)$ of the third order nonlinear current (Melrose 1978 Eqs. (10.105) and (10.125)). The matrix element for the decay of the O mode into the X mode and another transverse mode (which could be X or Alfvén wave) are calculated in appendix A. A typical growth rate for the decay of the O mode is (Eq. A4)

$$\Gamma \approx \frac{q^2}{m^2 c^3} \frac{\omega_p^6 \sin^4 \theta}{\gamma_p^3 \omega_B^2 \omega^4} W_k\tag{2}$$

where W_k is the wave intensity per unit interval of wave vectors k , q is the elementary charge, m is a mass of an electron, c is a velocity of light, ω_p is a plasma frequency, θ is an angle of propagation with respect to the magnetic field, ω_B is a cyclotron frequency, and γ_p is a Lorentz factor of the plasma motion.

Wave intensity W_k can be related to the observed flux F_ν : $W_k = \frac{1}{2\pi} F_\nu \left(\frac{d}{r}\right)^2$ where d/r is the ratio of the distance to the pulsar to the typical radius where transformations occur. If we parametrize the plasma density in terms of the Goldreich-Julian density $\omega_p^2 = 2\lambda_M \Omega \omega_B$ (where $\lambda_M \approx 10^6$ is a multiplicity factor that gives a number of secondary pairs per primary particle) then we can express the growth rate in terms of the observed flux and pulsar rotation frequency Ω

$$\Gamma \approx \frac{q^2}{m^2 c^3} \frac{\lambda_M^3 \Omega^3 \omega_{B,NS} \sin^4 \theta}{\gamma_p^3 \omega^4} \left(\frac{R_{NS}}{r}\right)^3 F_\nu \left(\frac{d}{r}\right)^2\tag{3}$$

where $\omega_{B,NS}$ is the surface cyclotron frequency and R_{NS} is the radius of the neutron star.

² Alfvén waves may be converted back to the X or O mode due to linear coupling in a strongly turbulent plasma. In any case the resulting waves will be unpolarized.

For a pulsar at $d = 1$ kpc, with rotation frequency $\Omega = 50$ rad/sec, average Lorentz factor of plasma $\gamma_p = 10$, observation frequency $\nu = 1$ GHz, typical interaction radius $r = 5 \times 10^6$ cm, and a peak flux of 1 Jy, we find the growth rate of the decay instability

$$\Gamma \approx 10^3 \sin^4 \theta \text{sec}^{-1} \quad (4)$$

This growth rate may be called marginal: efficient wave transformation occurs only for comparatively large angles of propagation $\theta \geq 0.1$ and it is quite sensitive to the choice of plasma parameters: changing the plasma density or the streaming Lorentz factor may make the growth time larger than the pulsar's period - making the wave transformation ineffective.

For a given intensity, the growth rate (3) decreases with radius $\propto r^{-3}$ (for a given observed flux growth rate $\propto r^{-5}$); thus, the most effective transformation occurs deep inside the pulsar magnetosphere. The effectiveness of the wave transformation in different pulsars depends mostly on the plasma production (λ_M and γ_p). Denser plasma streaming with a smaller Lorentz factor is more efficient for wave conversion. If the plasma production is independent of the pulsar period, then the wave transformation should be more efficient in the short period pulsars.

The frequency dependence of the wave transformation may also be complicated. On one hand, the efficiency of transformation (Eq. 3) is inversely proportional to the frequency, and the pulsar's spectral F_ν are usually power laws with index -1.6 (Lorimer et al. 1995). On the other hand, higher frequencies are believed to be emitted lower in the magnetosphere, so they acquire a larger angle with respect to the magnetic field in the interaction region; thus, they may have a larger decay rate. The lower frequencies, which are emitted at heights larger than $\sim$ ten stellar radii do not acquire large angles before they leave the region of effective wave transformation.

3. Conclusion

We showed that induced decay of the "upper" O mode into the unpolarized X mode in the inner parts of the pulsar magnetosphere may explain observed intensity dependent depolarization. When the intensity of the initial O mode is not high enough or the angle of propagation with respect to the magnetic field is small, the decay is weak and the original O mode is observed. When the intensity of the O mode is high and the wave is propagating at a considerable angle with respect to the magnetic field, it may completely decay into X modes (and backward propagating waves). Generally, the growth rate of the decay instability is marginal - variations in the plasma parameters and the structure of the magnetic field may make it either ineffective or extremely effective.

I would like to thank Anuj Parikh for his comments on the manuscript.

REFERENCES

- Arons J. & Barnard J.J. 1986, AJ, 302, 120
 Gedalin, M. E. & Machabeli G. Z. 1983, Sov. J. Plas. Phys., 9, 592
 Lorimer D.R. et al. 1995, MNRAS, 273, 411

- Lyutikov, M., Machabeli, G. & Blandford, R. 1999, ApJ, 512, 804
- Manchester R.N., Talor J.H. & Huguenin G.R. 1975, ApJ, 196, 83
- Melrose D.B. 1978, *Plasma astrophysics : nonthermal processes in diffuse magnetized plasmas*, New York, Gordon and Breach
- Xilouris K.M., Kramer M., Jessner A., Wielebinski R. 1995, A&A, 288, L17
- Zhelezniakov, V.V. 1996, *Radiation in astrophysical plasmas*, Dordrecht ; Boston Kluwer

A. Calculation of matrix element

Consider a decay of a high frequency ($\omega_O \gg \omega_P$) O mode into an X mode and another mode, unspecified at this point. Let the wave vectors and polarizations be

$$\begin{aligned} \mathbf{k}_0 &= k_0 \{\sin \theta, 0, \cos \theta\}, & \mathbf{e}_O &= \{\cos \theta, 0, \sin \theta\} \\ \mathbf{k}_X &= k_X \{\cos(\phi_X) \sin(\theta_X), \sin(\phi_X) \sin(\theta_X), \cos(\theta_X)\}, & \mathbf{e}_X &= \{\sin(\phi_X), \sin(\phi_X), 0\} \\ \mathbf{k}_2 &= k_2 \{\cos \phi_2 \sin(\theta_2), \sin \phi_2 \sin(\theta_2), \cos(\theta_2)\}, & \mathbf{e}_2 &= \{e_{2x}, e_{2y}, e_{2z}\} \end{aligned} \quad (\text{A1})$$

where θ and ϕ are polar coordinates in a frame aligned with the magnetic field, which is assumed to be in the z direction.

In a cold magnetized plasma the nonlinear third order currents (which determine the 3-wave interactions) are given by Melrose 1978 Eqs. (10.105) and (10.125). Two simplifications are possible in the case of strong magnetized electron-positron plasma. First, nonlinear current terms which are proportional to an odd power of the sign of the charge will cancel out. Secondly, we can make an expansion in powers $1/\omega_B$ and keep the lowest order. Under these assumption the matrix element is

$$V = \bar{S}_{ijl} e_{0,i}^* e_{X,j} e_{2,l} \approx -i \frac{e}{mc} \frac{\omega_p^2 \sin \theta}{\omega \omega_B} \sin(\phi_X - \phi_2) \quad (\text{A2})$$

where we assumed that the product waves propagate approximately along the field line, and wave 2 propagates backward.

Two conclusions may be made on analyzing Eq. (A2). First, for the decay to occur, the initial O mode should be obliquely propagating ($\theta \neq 0$). Secondly, the polarizations of the product wave are completely random with respect to the initial O mode and approximately orthogonal to each other. Thus, the decay of the linearly polarized O mode produces unpolarized waves. Microscopically, this is due to the fact that it is the z component of the initial O mode that couples to the product modes. A more physical description of the 3 wave interaction in pair plasma may be found in Machabeli & Rogava 1983.

Given the matrix element (A2) we can find the the probability of emission in the random phase approximation (which assumes that radiation is broadband) (Melrose 1978)

$$u = 4(2\pi)^7 V^2 \frac{\omega_X \omega_2}{\omega} \delta(\mathbf{k}_0 - \mathbf{k}_X - \mathbf{k}_2) \delta(\omega_0 - \omega_X - \omega_2). \quad (\text{A3})$$

Then the characteristic nonlinear decay time is

$$\Gamma \approx \frac{q^2}{m^2 c^3} \frac{\omega_p^6 \sin^4 \theta}{\gamma_p^3 \omega_B^2 \omega^4} W_k, \quad (\text{A4})$$

where we took into account that $\omega_2 = \omega_O - \omega_X \approx \frac{\omega_p^2 \sin^2 \theta}{\gamma_p^3 \omega}$.




Fig. 1.— Decay of O mode into X mode plus backward propagating X or Alfvén mode.